

Validated HPTLC method for simultaneous estimation of metoprolol succinate and ramipril in bulk drug and marketed formulation

Abstract

This paper describes a new, simple, precise, and accurate HPTLC method for simultaneous estimation of Metoprolol succinate and Ramipril as the bulk drug and in tablet dosage forms. Chromatographic separation of the drugs was performed on aluminum plates precoated with silica gel 60 F254 as the stationary phase and the solvent system consisted of Methanol: Toluene: Ethyl Acetate: Ammonia (2.5:3:5:0.7v/v/v/v). Densitometric evaluation of the separated zones was performed at 209 nm. The two drugs were satisfactorily resolved with R_F values 0.67, and 0.37 for Metoprolol Succinate and Ramipril, respectively. The accuracy and reliability of the method was assessed by evaluation of linearity (2000-12000 ng/spot for Metoprolol succinate and 200-1200 ng/spot for Ramipril), precision (intra-day RSD 0.471-1.036% and inter-day RSD 1.085-1.580% for Metoprolol Succinate and intra-day RSD 1.057-1.63% and inter-day RSD 1.024-1.746% for Ramipril), accuracy (98.95 ± 0.16 % for Metoprolol and 98.98 ± 0.41 % for Ramipril), and specificity, in accordance with ICH guidelines.

Keywords: Thin layer chromatography, densitometry, validation and quantification, Metoprolol Succinate, Ramipril.

Kamini Sethy^{1*}
Janhavi R. Rao²
K. Raja Rajeswari³
K.E.V. Nagoji⁴

Author Affiliations

^{1,3,4}Sri Venkateswara College of Pharmacy, Etcherla-532410, Srikakulam District, Andhra Pradesh, India.

²Poona College of Pharmacy, Bharati Vidyapeeth Deemed University, Erandwane, Pune-411038, Maharashtra, India.

*Corresponding Author

Kamini Sethy,
Sri Venkateswara College of Pharmacy, Etcherla-532410, Srikakulam District, Andhra Pradesh, India.

E-mail: kamini.tiki@gmail.com

Received on 22.06.2019

Accepted on 26.08.2019

1. Introduction

Metoprolol Succinate (Metoprolol) chemically is Bis [(2RS)-1-[4-(2-methoxyethyl) phenoxy]-3-[(1-methylethyl) amino] propan-2-ol] butanedioate (figure 1) is a cardio selective beta blocker. Metoprolol is used in the management of hypertension, angina pectoris, cardiac arrhythmias, myocardial infarction and heart failure. Metoprolol is also used in the management of hyperthyroidism and in the prophylactic treatment of migraine.

Ramipril chemically is 4-[2-(1-ethoxycarbonyl-3-phenyl-propyl)aminopropanoyl]-4-azabicyclooctane-3-carboxylic acid (figure 2) is a long acting angiotensin converting enzyme (ACE) inhibitor and it is a prodrug, which is hydrolysed after absorption to form the active metabolite ramiprilate which has a long elimination half-life, permitting once daily administration. Ramipril may be effective in indications such as severe essential hypertension and renal hypertension. It has also displayed beneficial effects in patients with moderate to severe congestive heart failure.




Figure 1: Structure of Metoprolol




Figure 2: Structure of Ramipril

Literature survey revealed that HPLC¹, HPTLC², LC-MS³ methods for estimation of Metoprolol individually and in combination with other drugs. Ramipril has been reported for the estimation as individual or in combination with other drugs in various analytical methods such as HPLC^{4,5,6,7,8,9,13}, HPTLC^{10,14}, LC-MS-MS¹¹, LC-MS¹², and spectrophotometric methods¹⁵ in bulk drug as well as in plasma, etc. Literature has revealed the analytical methods for simultaneous estimation of Metoprolol and Ramipril by RP-HPLC¹⁶ and spectrophotometry¹⁷. However, there is no analytical method reported till date for the simultaneous estimation of Metoprolol Succinate and Ramipril in a combined dosage formulation by HPTLC method. So this work was taken up for the development and validation by densitometric method which is advantageous over the existing methods in terms of sensitivity.

2. Experimental

Materials

Working standards of pharmaceutical grade Metoprolol (Batch no. 2148/009) and Ramipril (Batch no. 16043/01) obtained from Lupin Limited, Pune, India on dried basis as a gift sample. It was used without further purification. Commercial tablets of Metoprolol and Ramipril in a combined dosage form were purchased from the local market, brand name Starpress R XL-25 (Lupin). All chemicals and reagents (methanol, toluene, ethyl acetate, ammonia) used were of analytical grade and were purchased from Merck Chemicals, India.

Instrumentation

The samples were spotted in the form of bands of width 6 mm with a Camag 100 microlitre sample (Hamilton, Bonaduz, Switzerland) syringe on precoated silica gel aluminium plate 60 F - 254, (20 × 10 cm) with 250 µm thickness; E. Merck, Darmstadt, Germany, supplied by Anchrom Technologists, Mumbai) using a Camag Linomat IV applicator (Switzerland). The plates were prewashed by methanol and activated at 110°C for 5 min prior to chromatography. Then the chromatoplate was saturated with ammonia vapours for 30 min. A constant application rate of 0.1 µl/s was employed and space between two bands was kept at 6 mm. The slit dimension was kept at 5 × 0.45 mm and 10 mm/s scanning speed was employed. The monochromator bandwidth was set at 20 nm with K 320 cut off filter, each track was scanned thrice and baseline correction was used. The mobile phase


consisted of methanol: toluene: ethyl acetate: 30% ammonia (2.5: 3.0: 5.0, 0.7 v/v/v/v). 11.2 ml of mobile phase was used per chromatography. Linear ascending development was carried out in 20 × 10 cm twin trough glass chamber (Camag, Muttenz, Switzerland). Dimensions: length × width × height = 12 × 4.7 × 12.5 cm. It was saturated (lined on the two bigger sides with filter paper that had been soaked thoroughly with the mobile phase) and the chromatoplate development was carried out in dark with the mobile phase. The optimized chamber saturation time for mobile phase was 30 min at room temperature ($25^{\circ}\text{C} \pm 2$) at relative humidity of $60\% \pm 5$. The length of chromatogram run was 8 cm and approximately 20 min. Subsequent to the development, TLC plates were dried in a current of air with the help of an air dryer in wooden chamber with adequate ventilation. The flow of air in the laboratory was maintained unidirectional (laminar flow, towards exhaust). Densitometric scanning was performed on Camag TLC scanner III in the reflectance-absorbance mode at 209 nm for all measurements and operated by CATS software (V4.06, Camag). The source of radiation utilized was deuterium lamp emitting a continuous UV spectrum between 190 and 400 nm. Concentrations of the compound chromatographed were determined from the intensity of diffusely reflected light. Evaluation was via peak areas with linear regression.

Preparation of Standard Stock Solutions

20 mg of Metoprolol and 2 mg of Ramipril were accurately weighed and transferred to 10ml volumetric flask. Metoprolol and Ramipril were dissolved in 10ml of methanol to get Standard solutions of a concentration of 2 mg/ml of Metoprolol and 0.2 mg/ml of Ramipril. The standard solution was stored at $2-8^{\circ}\text{C}$, protected from light.

Optimization of the HPTLC method

The TLC procedure was optimized with a view to develop a simultaneous assay method for Metoprolol and Ramipril respectively. Various solvent systems like toluene: ethyl acetate: methanol, chloroform: methanol: ethyl acetate, toluene: ethyl acetate: methanol: ammonia was tried in different concentrations to separate and resolve spots of Metoprolol and Ramipril from their impurities and other excipients of formulations. Methanol: toluene: ethyl acetate: ammonia (2.5: 3.0: 5.0: 0.7 v/v/v/v) was found to result in the compact spot and best peak shape of Metoprolol and Ramipril. Metoprolol and Ramipril were satisfactorily resolved with R_f 0.67 ± 0.05 and 0.37 ± 0.02 respectively with acceptable resolution and peak shape (figure 3) at wavelength of 209 nm. In order to reduce the neckless effect TLC chamber was saturated for 20 min using saturation pads. The mobile phase was run upto a distance of 8 cm; which takes approximately 20 min for complete development of the TLC plate.



**Figure 3: HPTLC Densitogram of standard
Peak 1 (200 ng spot-1) of Ramipril (R_f 0.37 ± 0.02),
Peak 2 (2000 ng spot-1) of Metoprolol(R_f 0.67 ± 0.05),**

Validation of the method

Validation of the optimized TLC method was carried out with respect to the following parameters.

Linearity and range

From the mixed standard stock solution 2 mg/mL of Metoprolol and 0.2 mg/mL of Ramipril, 1 to 6 μ L solution spotted on TLC plate to obtain final concentration 2000-12000 ng/spot for Metoprolol and 200-1200 ng/spot for Ramipril. Linearity of the method was studied by injecting six concentrations of the drug each concentration was applied three times to the TLC plates. The plate was then developed using the previously described mobile phase and the peak areas were plotted against the corresponding concentrations to obtain the calibration curves.

Precision

The precision of the method was verified by repeatability and intermediate precision studies. Repeatability studies were performed by analysis of three different concentrations (2000, 6000, 10000 ng/spot for Metoprolol and 200, 600, 1000 ng/spot for Ramipril) of the drug six times on the same day. The intermediate precision of the method was checked by repeating studies on three different days.

Limit of detection and limit of quantitation

Limit of detection (LOD) and quantification (LOQ) represent the concentration of the analyte that would yield signal-to-noise ratios of 3 for LOD and 10 for LOQ, respectively. LOD and LOQ were determined by measuring the magnitude of analytical background by spotting a blank and calculating the signal-to-noise ratio for Metoprolol and Ramipril by spotting a series of solutions until the S/N ratio 3 for LOD and 10 for LOQ. To determine the LOD and LOQ, serial dilutions of mixed standard solution of Metoprolol and Ramipril were made from the standard stock solution in the range of 10–200 ng/spot. The samples were applied to TLC plate and the chromatograms were run and measured signal from the samples was compared with those of blank samples.

Robustness of the method

Following the introduction of small changes in the mobile phase composition (± 0.1 mL for each component), the effects on the results was examined. Mobile phases having different compositions, e.g. methanol: toluene: ethyl acetate: ammonia (2.6: 3: 5: 0.7 v/v/v/v), (2.5: 3.1: 5: 0.7 v/v/v/v), (2.5: 3: 5.1: 0.7v/v/v/v), (2.5: 3: 5: 0.8 v/v/v/v), were tried and chromatograms were run. The amount of mobile phase was varied over the range of ± 5 %. The plates were prewashed with methanol and activated at 60°C for 2, 5, and 7 min respectively prior to chromatography. The time from spotting to chromatography and from chromatography to scanning was varied from +10 min. The robustness of the method was determined at three different concentration levels 4000, 8000, 12000 ng/spot for Metoprolol and 400, 800, 1200 ng/spot for Ramipril.

Specificity

The specificity of the method was determined by analyzing standard drug and test samples. The spot for Metoprolol and Ramipril in the samples was confirmed by comparing the R_F and spectrum of the spot with that of a standard. The peak purity of Metoprolol and Ramipril was determined by comparing the spectrum at three different regions of the spot i.e. peak start (S), peak apex (M) and peak end (E).

Accuracy

Accuracy of the method was carried out by applying the method to drug sample (Metoprolol and Ramipril combination tablet) to which know amount of Metoprolol and Ramipril standard powder corresponding to 80, 100 and 120% of label claim had been added (standard addition), mixed and the powder was extracted and analyzed by running chromatogram in optimized mobile phase.

Analysis of a marketed formulation

To determine the content of Metoprolol and Ramipril in conventional tablet (Brand name: Starpress R XL25 Label claim: 25 mg Metoprolol and 2.5 mg Ramipril per tablet), ten tablets were weighed, their mean weight determined and finely powdered. The weight of the tablet triturate equivalent to 25 mg

Metoprolol and 2.5 mg Ramipril was transferred into a 25 mL volumetric flask containing 10-15 mL methanol, sonicated for 30 min and diluted to 25 mL with methanol. The resulting solution was centrifuged at 3000 rpm for 5 min and the drug content of the supernatant was determined (1000 and 100 µg/mL for Metoprolol and Ramipril respectively). 2µL of this solution (2000 and 200ng/spot for Metoprolol and Ramipril respectively) was applied to a TLC plate which was developed in optimized mobile phase. The analysis was repeated in triplicate. The possibility of excipient interference with the analysis was examined.

3. Results and Discussion

The results of validation studies on simultaneous estimation method developed for Metoprolol and Ramipril in the current study involving Methanol: toluene: ethyl acetate: ammonia (2.5: 3.0: 5.0: 0.7 v/v/v/v) as the mobile phase for TLC are given below.

Linearity

The drug response was linear ($r^2 = 0.997$ for Metoprolol and 0.999 for Ramipril) over the concentration range between 2000-12000 ng/spot for Metoprolol and 200-1200 ng/spot for Ramipril. The slope and intercept for Metoprolol and Ramipril were $1.284 (\pm 0.982)$, $1979 (\pm 1.25)$ and $2.947 (\pm 0.862)$ and $658 (\pm 1.06)$, respectively.

Precision

The results of the repeatability and intermediate precision experiments are shown in **Table 1**. The developed method was found to be precise as the RSD values for repeatability and intermediate precision studies were $< 2 \%$, respectively as recommended by ICH guidelines.

Table 1: Precision study for Metoprolol and Ramipril

Drug	Concentration ng per band	Intra-day (n = 3)		Inter-day (n = 3)	
		SD	RSD%	SD	RSD%
Metoprolol	60	14.28	1.040	18.15	1.326
	120	6.96	0.317	5.91	0.269
	180	24.83	0.865	32.73	1.141
Ramipril	60	56.65	1.904	49.86	1.708
	120	33.97	0.671	31.33	0.618
	180	40.51	0.627	41.03	0.635

LOD and LOQ

Signal-to-noise ratios of 3: 1 and 10: 1 were obtained for the LOD and LOQ respectively. The LOD and LOQ were found to be 50 ng/spot and 100 ng/spot for Metoprolol and 50 ng/spot and 150 ng/spot for Ramipril, respectively.

The standard deviation of peak areas was calculated for each parameter and the % RSD was found to be less than 2 %. The low values of the % RSD, as shown in **Table 2** indicated robustness of the method.

Table 2: Robustness Testing of Metoprolol and Ramipril

Parameters	Metoprolol		Ramipril	
	SD	%RSD*	SD	%RSD*
Mobile phase composition (± 0.1 ml)	10.42	1.235	10.42	1.235
Amount of mobile phase (± 0.5 %)	20.14	1.018	20.14	1.018
Time from spotting to chromatography (± 20 min)	15.36	0.942	15.36	0.942
Time from chromatography to scanning (± 20 min)	20.10	1.085	20.10	1.085

*n=3

Specificity

The peak purity of Metoprolol and Ramipril was assessed by comparing their respective spectra at the peak start, apex, and peak end positions of the spot, i.e., $r(S, M)=0.998$ and $r(M, E)=0.999$. A good correlation ($r=0.9997$) was also obtained between the standard and sample spectra of Metoprolol and Ramipril, respectively. Also, excipients from formulation were not interfering with the assay.

Recovery Studies

As shown from the data in Table 3 good recoveries of the Thiocolchicoside and Aceclofenac in the range from 98.32 to 99.45 % were obtained at various added concentrations. The average recovery of three levels (nine determinations) for Metoprolol and Ramipril were 98.95 % and 98.98 % respectively.

Table 3: Recovery studies of Metoprolol and Ramipril

Label claim (mg/tablet)	Amount Added (%)	Total amount (mg)	Amount* recovered (mg \pm % RSD)	Recovery (%)
Metoprolol 25	80 (20mg)	45	44.65 \pm 0.222	99.22
	100 (25mg)	50	49.16 \pm 0.154	98.32
	120 (30mg)	55	54.63 \pm 0.130	99.32
Ramipril 2.5	80 (2mg)	4.5	4.46 \pm 0.526	99.11
	100 (2.5mg)	5.0	4.92 \pm 0.360	98.4
	120 (3mg)	5.5	5.47 \pm 0.344	99.45

*n =3

Analysis of a formulation

Experimental results of the amount of Metoprolol and Ramipril in tablets, expressed as a percentage of label claims were in good agreement with the label claims thereby suggesting that there is no interference from any of the excipients, which are normally present in tablets. The average drug content was found to be 99.52 % and 99.6 % for Metoprolol and Ramipril, respectively, using the proposed procedures and the results are summarized in Table 4.

Table 4: Results from assay of Metoprolol, Ramipril in Starpress R XL 25 tablet

Component	Label claim(mg)	Amount found Mg \pm SD, n = 6	Percentage of label claim (\pm SD)
Metoprolol	25	24.88 \pm 35.84	99.52% \pm 15.15
Ramipril	2.5	2.49 \pm 18.87	99.6% \pm 10.21

4. Conclusion

The developed TLC technique is precise, specific and accurate. Statistical analysis proves that the method is suitable for the analysis of Metoprolol and Ramipril as bulk drug and in pharmaceutical formulation without any interference from the excipients. It may be extended to study the degradation kinetics of Metoprolol and Ramipril and also for its estimation in plasma and other biological fluids. The proposed TLC method is less expensive, simpler, rapid, and more flexible than HPLC.

Acknowledgement

The authors would like to thank, Lupin Limited, Pune, India for providing gift samples of standard Metoprolol Tartarate and Ramipril. The authors would like to thank, Dr. KR. Mahadik, Principal, Poona College of Pharmacy, Pune, Bharti Vidyapeeth Deemed University, India for providing necessary facilities to carry out the work.

References

1. Mitesh D Phale, Purnima D Hamrapurkar, A validated and simplified RP-HPLC of Metoprolol succinate from bulk drugs, Asian J. Research Chem.(2): April- June, 2009
2. Rajendra Kakde, Nilesh Bawane, High-performance thin-layer chromatographic method for simultaneous analysis of metoprolol succinate and amlodipine besylate in pharmaceutical preparations , Journal of Planar Chromatography, Volume 22, Number 2/ April 2009 pages 115-119
3. P. Senthamil Selvan and T.K. Pal, Chromatography-tandem mass spectrometry method for the simultaneous quantitation of metoprolol succinate and simvastatin in human plasma. 1.Volume 49, Issue 3, 5 April 2009, Pages 780-785.
4. F. Belal, I.A. Al-Zaagi, E.A. Gadkariem and M. A. Abounassif, A stability-indicating LC method for the simultaneous determination of ramipril and hydrochlorothiazide in dosage forms. Journal of Pharmaceutical and Biomedical Analysis, Volume 24, Issue 3, January 2001, Pages 335-342.
5. K. Srinivasa Rao, K. Srinivas, RP-HPLC method for the determination of Losartan Potassium and Ramipril in combined dosage form. Indian J. Pharm. Sci., 2010, 72(1): 108-111.
6. Lincy Joseph, Mathew George, Venkata Ranga Rao B, Simultaneous estimation of Atorvastatin and Ramipril by RP-HPLC and spectroscopy. Pak. J. Pharm. Sci., Vol. 21, No.3 July 2008, pp.282-284.
7. Hiral J. Panchal, Bhanubhai N. Suhagia, Natvarlal J. Patel, Ishwarsinh S. Rathod, Bhavesh H.Patel, Simultaneous estimation of Atorvastatin Calcium, Ramipril and Aspirin in capsule dosage form by RP-HPLC, Chromatographia 2009, January (No. ½)
8. V P Kurade, MG Pai, R Gude, RP-HPLC estimation of Ramipril and Telmisartan in tablets, Indian Journal of Pharmaceutical Sciences, 2009., Vol. 71, Issue2, page148-151
9. M.M. Baing, V.V. Vaidya, R.T. Sane, S.N. Menon, K. Dalvi, Simultaneous RP-HPLC determination of Losartan Potassium, Ramipril, and Hydrochlorothiazide in pharmaceutical preparations, Chromatographia 2006, 64, September (No. 5/6)
10. Hiral J. Panchal, Bhanubhai N. Suhagia, Natvarlal J. Patel, Simultaneous HPTLC analysis of Atorvastatin Calcium, Ramipril and Aspirin in a capsule dosage form. Journal of Planar Chromatography, Volume 22, Number 4/ August 2009
11. Bo Yuan, Xuan Wang, Fenge Zhang, Jun Jia, Fangling Tang, Simultaneous Determination of Ramipril and its active metabolite Ramiprilat in human plasma by LC-MS-MS. Chromatographia, 2008., Volume 68, pages 533-539
12. K. Veeran Gowda, Uttam Mandal, P. Senthamil Selvan, W.D. Sam Solomon, Animesh Ghosh, Amlan Kanti Sarkar, Sangita Agarwal, T. Nageswar Rao and Tapan Kumar Pal, Liquid chromatography tandem mass spectrometry method for simultaneous determination of metoprolol tartrate and ramipril in human plasma. Journal of Chromatography B , Volume 858, Issues 1-2, 15 October 2007, Pages 13-21

13. Bilal YILMAZ Determination of Ramipril in pharmaceutical preparations by high-performance liquid chromatography. International Journal of Pharmaceutical Sciences Review and Research , Volume 1, Issue 1, March – April 2010; Page 39
14. Laxman V. Potale, Mrinalini C. Damle, Amol S. Khodke¹ and K.G. Bothara A validated stability indicating HPTLC method for simultaneous estimation of Ramipril and Telmisartan. International Journal of Pharmaceutical Sciences Review and Research Volume 2, Issue 2, May – June 2010; Article 007
15. Priyanka R Patil, Sachin U Rakesh, Pandurang N Dhabale, Kishor B Burade. Simultaneous Estimation of Ramipril and Amlodipine by UV Spectrophotometric Method, Research J. Pharm. and Tech.2 (2): April.-June. 2009,
16. R.J. Chandra Bose; G. Sivanseyal; Krishnakumar D, Validated Rp-Hplc Method for the Simultaneous Estimation of Ramipril and Metoprolol Tartrate in Bulk and Tablet Dosage Form, Asian Journal of Biochemical and Pharmaceutical Research, Volume 1, Issue 2, p.171-177.
17. K. Suresh Kumar, R. Ravikumar, A. Rajasekaran, simultaneous spectrophotometric determination of Metoprolol tartarate and Ramipril; Digest Journal of Nanomaterials and Biostructures; Vol. 5, No 1, March 2010, p. 173-176